

# Fermi-Operator Expansions for Linear Scaling Electronic Structure Calculations


---

Aiichiro Nakano

*Collaboratory for Advanced Computing & Simulations  
Department of Computer Science  
Department of Physics & Astronomy  
Department of Chemical Engineering & Materials Science  
Department of Quantitative & Computational Biology  
University of Southern California*

Email: [anakano@usc.edu](mailto:anakano@usc.edu)


$O(N)$  sparse matrix representation  
Simple & generalizable → use it



# Fermi Operator

- Fermi operator

$$F(\hat{H}) = \frac{2}{\exp\left(\frac{\hat{H}-\mu}{k_B T}\right) + 1}$$



- Projection to the occupied subspace

$$|\psi_{\text{proj}}\rangle = F(\hat{H}) |\psi\rangle$$

- The expectation value of any operator  $A$  is obtained by

$$\langle \hat{A} \rangle = \text{tr}[\hat{A} F(\hat{H})]$$

- Widely used in  $O(N)$  electronic structure calculations ( $N$  = number of electrons) through its sparse representation

cf.  $O(N^3)$  way



$$\hat{H}|n\rangle = \varepsilon_n |n\rangle$$

$$\langle \hat{A} \rangle = \sum_n \frac{2}{\exp\left(\frac{\varepsilon_n - \mu}{k_B T}\right) + 1} \langle n | \hat{A} | n \rangle$$

# Fermi-Operator Approximations

## Chebyshev polynomial

$$T_n(\cos\theta) = \cos(n\theta)$$




## Rational

$$F(\hat{H}) \cong \sum_{\nu=1}^M \frac{R_\nu}{\hat{H} - z_\nu}$$

$$(\hat{H} - z_\nu) |\psi_{\text{out}}^\nu\rangle \cong R_\nu |\psi_{\text{in}}\rangle$$

$$F(\hat{H}) |\psi_{\text{in}}\rangle \cong \sum_{\nu=1}^M |\psi_{\text{out}}^\nu\rangle$$



See note on Fermi-operator expansion

# Rational Fermi-Operator Expansion

$$\begin{aligned} f(z) &= \frac{1}{\exp(z) + 1} & e^z &= \lim_{n \rightarrow \infty} \left(1 + \frac{z}{n}\right)^n \\ &\cong \frac{1}{\left(1 + \frac{z}{2M}\right)^{2M}} \\ &\cong \sum_{\nu=0}^{2M-1} \frac{R_\nu}{z - z_\nu} \end{aligned}$$


**Poles**

$$z_\nu = 2M \left( \exp \left( i \frac{(2\nu + 1)\pi}{2M} \right) - 1 \right)$$

**Residues**

$$R_\nu = - \exp \left( i \frac{(2\nu + 1)\pi}{2M} \right)$$

$$(\nu = 0, \dots, 2M - 1)$$



D. M. C. Nicholson *et al.*, *Phys. Rev. B* **50**, 14686 ('94); A. P. Horsfield *et al.*, *Phys. Rev. B* **53**, 12694 ('96); L. Lin *et al.*, *J. Phys. Condens. Matter* **25**, 1295501 ('13)

See note on Fermi-operator expansion

# $O(N)$ Fermi Operator Expansion

- Truncated expansion of Fermi-operator by Chebyshev polynomial  $\{T_p\}$

$$F(\hat{H}) \cong \sum_{p=0}^P c_p T_p(\hat{H})$$

- $O(N)$  algorithm

prepare a basis set of size  $O(N)$

(let the size be  $N$  for simplicity)

for  $l = 1, N$

let an  $N$ -dimensional unit vector be  $|e_l\rangle = \begin{bmatrix} 0 \\ \vdots \\ 1 \\ \vdots \\ 0 \end{bmatrix}_l$  *l-th atomic-site orbital*  
recursively construct the  $l^{\text{th}}$  column of  
matrix  $T_p$ ,  $|t_l^p\rangle$ , keeping only  $O(1)$

off-diagonal elements\* (*cf.* quantum nearsightedness<sup>#</sup>)

$$\left\{ \begin{array}{l} |t_l^0\rangle = |e_l\rangle \\ |t_l^1\rangle = \hat{H}|e_l\rangle \\ |t_l^{p+1}\rangle = 2\hat{H}|t_l^p\rangle - |t_l^{p-1}\rangle \end{array} \right. \quad \text{cf. Legendre polynomial by recursion}$$

build a sparse representation of the  $l^{\text{th}}$  column of  $F$  as

$$|f_l\rangle = \sum_{p=0}^P c_p |t_l^p\rangle$$

\*Six degrees of separation

#W. Kohn, *Phys. Rev. Lett.* **76**, 3168 ('96)