


Density Functional Theory (DFT)

Aiichiro Nakano

*Collaboratory for Advanced Computing & Simulations
Depts. of Computer Science, Physics & Astronomy, Chemical
Engineering & Materials Science, and Biological Sciences
University of Southern California*

Email: anakano@usc.edu

**How to determine the electronic ground state
in Born-Oppenheimer molecular dynamics (BOMD)?**



Hohenberg-Kohn Theorem

Consider a system of N electrons in an external potential $v(\mathbf{r})$.

Theorem 1

The ground-state density $\rho(\mathbf{r})$ & the external potential $\{v(\mathbf{r}) + c\}$ (c is a constant) are bijective functional, *i.e.*, one-to-one correspondence (see the lecture note on DFT for a proof).

Theorem 2

Any property of the many-electron ground state $|\Psi\rangle$, including the ground-state energy, $E = \langle\Psi|H|\Psi\rangle$ (H is the Hamiltonian operator), is a functional of $\rho(\mathbf{r})$.

P. Hohenberg & W. Kohn, "Inhomogeneous electron gas," *Phys. Rev.* **136**, B864 ('64)

- See lecture notes on (1) functional derivative & (2) density functional theory

Energy Functional


Definition: Exchange-correlation (xc) functional

$$E[\rho(\mathbf{r})] = T_s[\rho(\mathbf{r})] + \int d\mathbf{r} v(\mathbf{r})\rho(\mathbf{r}) + \frac{1}{2} \int d\mathbf{r} d\mathbf{r}' \frac{\rho(\mathbf{r})\rho(\mathbf{r}')}{|\mathbf{r} - \mathbf{r}'|} + E_{xc}[\rho(\mathbf{r})]$$

Kinetic energy of
non-interacting
electrons

Hartree energy (mean-
field approximation to
the electron-electron
interaction energy)

Exchange-correlation
energy



Kohn-Sham Equation

- The many-electron ground state is obtained by solving a set of one-electron Schrödinger equations called Kohn-Sham (KS) equations

$$\left[-\frac{\hbar^2}{2m} \frac{\partial^2}{\partial \mathbf{r}_i^2} + v_{\text{KS}}(\mathbf{r}) \right] \psi_n(\mathbf{r}) = \epsilon_n \psi_n(\mathbf{r})$$

KS wave function

KS energy

- KS potential**

$$v_{\text{KS}} = v(\mathbf{r}) + \int d\mathbf{r}' \frac{e^2 \rho(\mathbf{r}')}{|\mathbf{r} - \mathbf{r}'|} + v_{\text{xc}}(\mathbf{r})$$

exchange-correlation (xc) potential

$$\rho(\mathbf{r}) = \sum_n \Theta(\mu - \epsilon_n) |\psi_n(\mathbf{r})|^2$$

step function chemical potential

$$v_{\text{xc}}(\mathbf{r}) \equiv \frac{\delta E_{\text{xc}}}{\delta \rho(\mathbf{r})}$$

$$N = \sum_n \Theta(\mu - \epsilon_n)$$

- See the lecture note on density functional theory for derivation

W. Kohn & L. J. Sham, "Self-consistent equations including exchange and correlation effects," *Phys. Rev.* **140**, A1133 ('65)

Extensions of DFT

- **Finite-temperature DFT**

N. D. Mermin, “Thermal properties of the inhomogeneous electron gas,”
Phys. Rev. **137**, A1441 ('65)

- **Current DFT in magnetic fields: $\rho(\mathbf{r})$ & $\mathbf{j}(\mathbf{r})$**

M. Ferconi & G. Vignale, “Current density functional theory of quantum dots in magnetic fields,” *Phys. Rev. B.* **50**, 14722 ('94)

- **Superconducting-gap DFT: $\Delta(\mathbf{r}) = \langle \psi_{\uparrow}(\mathbf{r})\psi_{\downarrow}(\mathbf{r}) \rangle$ & $\rho(\mathbf{r}) = \sum_{\sigma} \langle \psi_{\sigma}^{\dagger}(\mathbf{r})\psi_{\sigma}(\mathbf{r}) \rangle$**

L. N. Oliveira *et al.*, “Density functional theory for superconductors,”
Phys. Rev. Lett. **60**, 2430 ('88)

- **For the electronic current operator, see the lecture note on quantum dynamical computation of electronic conductivity**